

# WEIGHT MANAGEMENT

**Zinc** Deficiency of zinc reduces leptin, a beneficial hormone that regulates appetite, which is reversed by zinc repletion.<sup>10,37</sup>

**Asparagine** This amino acid increases insulin sensitivity which helps the body store energy in muscle instead of storing it as body fat.<sup>1,2</sup>

**Biotin** Boosts metabolism by improving glycemic control (stabilizes blood sugar) and lowering insulin, a hormone that promotes fat formation.<sup>3,4,5</sup>

**Carnitine** Carries fatty acids into the cell so they can be burned for fuel; Helps reduce visceral adiposity (belly fat).<sup>6,7</sup>

**Calcium** Inhibits the formation of fat cells; Also helps oxidize (burn) fat cells.<sup>8,9,10</sup>

**Lipoic Acid** Improves glucose uptake into cells, which helps a person burn carbohydrates more efficiently.<sup>11,12,13</sup>

**Chromium** Makes the body more sensitive to insulin, helping to reduce body fat and increase lean muscle.<sup>14,15,16,27,28,4</sup>

**Vitamin B5** Taking B5 lowers body weight by activating lipoprotein lipase, an enzyme that burns fat cells. One study linked B5 supplementation to less hunger when dieting.<sup>17,18</sup>

**Magnesium** Low magnesium in cells impairs a person's ability to use glucose for fuel, instead storing it as fat; Correcting a magnesium deficiency stimulates metabolism by increasing insulin sensitivity. Magnesium may also inhibit fat absorption.<sup>19,20,21</sup>

**Glutamine** Reduces fat mass by improving glucose uptake into muscle.<sup>22,23</sup>

**Cysteine** Supplementation with this antioxidant reduced body fat in obese patients.<sup>24</sup>

**Inositol** Supplementation may increase adiponectin levels.<sup>25</sup>

**Vitamin B3 (Niacin)** Treatment with B3 increases adiponectin, a weight-loss hormone secreted by fat cells; Niacin-bound chromium supplements helped reduced body weight in clinical trials.<sup>26,27,28</sup>

**Vitamin A** Enhances expression of genes that reduce a person's tendency to store food as fat; Reduces the size of fat cells.<sup>10,29,30</sup>

**Vitamin E** Inhibits pre-fat cells from changing into mature fat cells, thus reducing body fat.<sup>10,31,32</sup>

**Vitamin D** Deficiency strongly linked to poor metabolism of carbohydrates; Genes that are regulated by vitamin D may alter the way fat cells form in some people.<sup>8,33,34</sup>

**Vitamin K** Poor vitamin K status linked to excess fat tissue; Vitamin K helps metabolize sugars.<sup>35,36</sup>

## REFERENCES

- <sup>1</sup>Lancha A, Poortmans J, Pereira L. The effect of 5 days of aspartate and asparagine supplementation on glucose transport activity in rat muscle. *Cell Biochem Funct* 2009;8:552-557.
- <sup>2</sup>Marquezi M, Roschel H et al. Effect of aspartate and asparagine supplementation on fatigue determinants in intense exercise. *Int J Sport Nutr Exer Metab* 2003;13:65-75.
- <sup>3</sup>Larrieta E, de la Vega-Monroy M, Vital P et al. Effects of biotin deficiency on pancreatic islet morphology, insulin sensitivity and glucose homeostasis. *J Nutr Biochem* 2012;4:392-399.
- <sup>4</sup>Albarracin C, Fuqua B, Evans J et al. Chromium picolinate and biotin combination improves glucose metabolism in treated, uncontrolled overweight to obese patients with type 2 diabetes. *Diabetes Metab Res* 2008;1:41-51.
- <sup>5</sup>Monograph on Biotin. *Altern Med Rev* 2007;12:73-78.
- <sup>6</sup>Bernard A, Rigault C, Mazue F et al. L-carnitine supplementation and physical exercise restore age-associated decline in some mitochondrial functions in the rat. *J Gerontol A Biol Sci Med Sci* 2008;10:1027-1033.
- <sup>7</sup>Galloway S, Craig T, Cleland S. Effects of oral L: -carnitine supplementation on insulin sensitivity indices in response to glucose feeding in lean and overweight/obese males. *Amino Acids* 2011;2:507-515.
- <sup>8</sup>Rosenblum J, Castro V, Moore C et al. Calcium and vitamin D supplementation is associated with decreased abdominal visceral adipose tissue in overweight and obese adults. *Am J Clin Nutr* 2012;1:101-108.
- <sup>9</sup>Zemal M. Role of calcium and dairy products in energy partitioning and weight management. *Am J Clin Nutr* 2004;79(Suppl):S907-S912.
- <sup>10</sup>Garcia O, Long K, Rosado J. Impact of micronutrient deficiencies on obesity. *Nutr Rev* 2009;10:559-572.
- <sup>11</sup>Teachey M, Taylor Z, Maier T et al. Interactions of conjugated linoleic acid and lipoic acid on insulin action in the obese Zucker rat. *Metabolism* 2003;9:1167-1174.
- <sup>12</sup>Zhang Y, Han P, Wu N et al. Amelioration of Lipid Abnormalities by  $\alpha$ -Lipoic acid Through Antioxidative and Anti-Inflammatory Effects. *Obesity* 2011;8:1647-1653.
- <sup>13</sup>Ansar H, Mazloom Z, Kazemi F et al. Effect of alpha-lipoic acid on blood glucose, insulin resistance and glutathione peroxidase of type 2 diabetic patients. *Saudi Med J* 2011;6:584-588.
- <sup>14</sup>Kim C, Kim B, Park K et al. Effects of short-term chromium supplementation on insulin sensitivity and body composition in overweight children: randomized, double-blind, placebo-controlled study. *J Nutr Biochem* 2011;11:1030-1034.
- <sup>15</sup>Lau F, Bagchi M, Sen C et al. Nutrigenomic basis of beneficial effects of chromium(III) on obesity and diabetes. *Mol Cell Biochem* 2008;1-2:1-10.
- <sup>16</sup>Cefalu W, Rood J, Pinsonat P et al. Characterization of the metabolic and physiologic response to chromium supplementation in subjects with type 2 diabetes mellitus. *Metabolism* 2010;5:755-762.
- <sup>17</sup>Naruta E, Buko V. Hypolipidemic effect of pantothenic acid derivatives in mice with hypothalamic obesity induced by aurothioglucose. *Exp Toxicol Pathol* 2001;5:393-398.
- <sup>18</sup>Leung L. Pantothenic acid as a weight-reducing agent: fasting without hunger, weakness and ketosis. *Med Hypotheses* 1995;5:403-405.
- <sup>19</sup>Takaya J, Higashino H, Kobayashi Y. Intracellular magnesium and insulin resistance. *Magnes Res* 2004;2:126-36.
- <sup>20</sup>Kishimoto Y, Tani M, Uto-Kondo H et al. Effects of magnesium on postprandial serum lipid responses in healthy human subjects. *Br J Nutr* 2010;4:469-472.
- <sup>21</sup>Lima M, Cruz T, Rodrigues L et al. Serum and intracellular magnesium deficiency in patients with metabolic syndrome—evidences for its relation to insulin resistance. *Diabetes Res Clin Pract* 2009;2:257-262.
- <sup>22</sup>Greenfield J, Farooqi I, Keogh J et al. Oral glutamine increases circulating glucagon-like peptide 1, glucagon, and insulin concentrations in lean, obese, and type 2 diabetic subjects. *Am J Clin Nutr* 2009;1:106-113.
- <sup>23</sup>Prada P, Hirabara S, de Souza C et al. L-glutamine supplementation induces insulin resistance in adipose tissue and improves insulin signalling in liver and muscle of rats with diet-induced obesity. *Diabetologia* 2007;9:1949-59.
- <sup>24</sup>Hildebrandt W, Hamman A, Krakowski-Roosen H et al. Effect of thiol antioxidant on body fat and insulin reactivity. *J Mol Med* 2004;5:336-344.
- <sup>25</sup>Corrado F, D'anna R, Di Vieste G et al. The effect of myoinositol supplementation on insulin resistance in patients with gestational diabetes. *Diabet Med* 2011;8:972-975.
- <sup>26</sup>Westphal S, Borucki K, Taneva E et al. Adipokines and treatment with niacin. *Metabolism* 2006;10:1283-1285.
- <sup>27</sup>Rink C, Roy S, Khanna S et al. Transcriptome of the subcutaneous adipose tissue in response to oral supplementation of type 2 Leprdb obese diabetic mice with niacin-bound chromium. *Physiol Genomics* 2006;3:370-379.
- <sup>28</sup>Preuss H, Bagchi D, Bagchi M et al. Effects of a natural extract of (-)-hydroxycitric acid (HCA-SX) and a combination of HCA-SX plus niacin-bound chromium and Gymnema sylvestre extract on weight loss. *Diabetes Obes Metab* 2004;3:171-180.
- <sup>29</sup>Kameji H, Mochizuki K, Myoshi N et al.  $\beta$ -Carotene accumulation in 3T3-L1 adipocytes inhibits the elevation of reactive oxygen species and the suppression of genes related to insulin sensitivity induced by tumor necrosis factor- $\alpha$ . *Nutrition* 2010;11-12:1151-1156.
- <sup>30</sup>Ribot J, Felipe F, Bonet M et al. Changes in adiposity in response to vitamin A status correlate with changes of PPAR gamma 2 expression. *Obes Res* 2001;9:500-509.
- <sup>31</sup>Ohrvall M, Tengblad S, Vessby B. Lower tocopherol serum levels in subjects with abdominal adiposity. *J Intern Med* 1993;234:53-60.
- <sup>32</sup>Uto-Kondo H, Ohmori R, Kiyose C et al. Tocotrienol Suppresses Adipocyte Differentiation and Akt Phosphorylation in 3T3-L1 Preadipocytes. *J Nutr* 2009;1:51-57.
- <sup>33</sup>Bailey R, Cooper J, Zeitels K et al. Association of the vitamin D metabolism gene CYP27B1 with type I diabetes. *Diabetes* 2007;10:2616-2621.
- <sup>34</sup>Ochs-Balcom H, Chennamaneni R, Millen A et al. Vitamin D receptor gene polymorphisms are associated with adiposity phenotypes. *Am J Clin Nutr* 2011;1:5-10.
- <sup>35</sup>Yoshida M, Jacques P, Meigs J et al. Effect of vitamin K supplementation on insulin resistance in older men and women. *Diabetes Care* 2008;11:2092-2096.
- <sup>36</sup>Shea M, Booth S, Gundberg C. et al. Adulthood obesity is positively associated with adipose tissue concentrations of vitamin K and inversely associated with circulating indicators of vitamin K status in men and women. *J Nutr* 2010;5:1029-1034.
- <sup>37</sup>Jansen J et al. Zinc and diabetes—clinical links and molecular mechanisms. *J Nutr Biochem* 2009;6:399-417.